

Preparation and Shipping Procedures

Ship samples packed in 8-12 kg of dry ice* to:

Metabolon Sample Acceptance 617 Davis Drive, Suite 400 Morrisville, NC 27560

Phone: +1-919-572-1711

Ship domestic overnight Monday - Wednesday. Ship international via World Courier, Monday or Tuesday Note: Our receiving hours are 09:00 to 16:00 ET.

Shipment Details

A completed digital manifest is required for each shipment, and a template will be provided by your Metabolon Client Success Project Manager. Please include a hard copy of the completed manifest inside your sample shipment but outside of the Styrofoam container.

Prior to shipping samples, email an electronic copy of the completed sample manifest in Excel format to: samplemanager@metabolon.com. Please include your assigned project code in the subject line. Please do not ship samples until your project code has been assigned.

Remember to include a hard copy of the completed manifest with your sample shipment

Completing Your Digital Manifest:

- Must include the project code on the cover page and in the subject line of all email correspondence.
- Must include all available sample information for each sample shipped, including any preservatives used in the collection/storage process.
- Must NOT include personally identifying information for human samples.

Planning

- Complying with required sample amounts and consistent handling is critical to study quality (e.g., collection technique, time of sampling, time to freezer, freeze/thaw, time of aliquoting).
- Please notify your Metabolon Study Director of any preservatives that have been used in the collection process, novel sample types, and samples containing less than minimum required volumes or masses.
- The sample disposition date will be provided in 90 days post project completion.

Aliquot and Freeze

- Minimize or eliminate the time samples are in a nonfrozen state.
- Transfer required volume or mass of sample into pre-labeled and pre-chilled tubes, then freeze.
- ► Flash-freeze the sample vials in liquid nitrogen or immediately store them in a -80 °C freezer.

Acceptable Sample Tubes

- The Metabolon Study Success Sample Handling Kit will provide you with Metabolon Standard Preferred 2D barcoded sample collection tubes specifically intended for your study.
- For information on other barcoded tubes that may be acceptable, please contact your Metabolon Client Success Project Manager.
- Scan the sample barcode into the supplied digital manifest and provide additional requested metadata. The minimum required information fields within the digital manifest include the following: "Unique Tube Label ID", "Client Matrix", "Sample Amount", "Sample Amount Units" "Sample Container ID" and "Sample Container Position".

Collection Guidelines

Blood Samples Acceptable sample types:

Plasma:

Collect whole blood in anti-coagulant tubes following the manufacturer's processing instructions.

Plasma Anticoagulant Guidelines:

- Best results: EDTA (K2, K3, Na; avoid Li).
- For global metabolomics studies, avoid citrate.
- For all studies, never include multiple anticoagulant sample types in the same experiment.

Serum:

 Collect whole blood in serum separator tubes and follow the tube manufacturer's processing instructions.

Whole blood:

 Coagulation must be prevented, preferably by the addition of EDTA.

Fecal Samples

Frozen:

Do not lyophilize samples unless instructed to by Metabolon.

OMNImet•GUT[™] Fecal Samples:

- Gather all individual donor samples at your facility before shipping to Metabolon. We will not accept samples direct from individual donors.
- Store samples at -80°C as soon as possible.
 - Room temperature stability is validated for 4 days (Global Metabolomics Platform) or 7 days (Short-Chain Fatty Acid Assay).
 - ► Store at -20°C for up to 1 week if -80°C is unavailable. Avoid exposure to temperatures above 30°C (86°F).
- Provide exposure to temperatures above 50 C (66 F) Provide donors with ice packs if such exposure is likely prior to arrival at your facility.
- Ship OMNImet-GUT tubes to Metabolon in their entirety. If you must aliquot samples, please consult with your Project Coordinator first.
- Follow standard shipping procedures (page 1) once all donor samples are gathered.

Tissue Samples

For solid tissues (e.g., biopsy material), the amount of tissue per sample can vary depending upon study objectives and tissue type. The minimum amount requirement is 50 mg of sample.

Skin Samples

Discuss with your Metabolon representative.

Cell Samples (Eukaryotic and Bacterial)

- Pellet cells and remove supernatant.
- Provide an accurate cell count for each sample.

Plant Material

 Preferably, grind fresh plant material to a powder with a mortar andpestle under liquid nitrogen. The powdered material should be lyophilized, if possible.

Dried Blood Spot Cards

- ► Submit dried Whatman[™] 903 Protein Saver DBS cards, including 2 blank cards per study in separate zip-top bags (Whatman# 10548232, VWR# 89027-022) containing silica gel desiccant pack.
- Store bags together in a secondary vessel containing a humidity indicator card.
- ▶ For maximum stability, we recommend -80°C storage.
- Samples should be shipped on dry ice with a humidity indicator card and silica gel desiccant included with the shipment.

Biohazardous Materials

 BSL2 solids and liquids are acceptable. Notify Metabolon of the pathogen involved.

Liquid Sampling Procedure

(Blood, Urine, cell culture media, or CSF)

- 1. Pipet required volume per sample type as defined in the Table.
- 2. Immediately dispense pipetted liquid into the chilled polypropylene tube and flash-freeze.
- 3. Store samples at -80 °C until shipment.
- 4. Volumes below the minimum requirement should be discussed with Study Director.

Solid Sampling Procedure

(Feces, Tissue, or Plant)

- Weigh the required mass per sample as defined in the Table. (Do not include any homogenization materials such as metal or ceramic beads, garnet, etc.)
- 2. Transfer weighed sample into the chilled polypropylene tube and flash-freeze.
- 3. Store samples at -80 °C until shipment.
- 4. Mass below the minimum requirement should be discussed with your Study Director

Specimen Type	Min. Amount
Blood	150 to 300 μL
Urine	150 to 300 μL
Cerebrospinal Fluid (CSF)	150 to 300 μL
Cell Culture Media	150 to 300 µL
Cells (Packed cell volume)	≥100 µL
Fecal	100 mg
Tissue	50 mg
Plant Material (Dry Weight)	50 to 100 mg
Dried Blood Spots (diameter)	8-12 mm (x2)
Sebutape or D-Squame Discuss with your Metal	bolon representative

Sample Amount Requirements

- Sample amounts per assay are defined in the above table. Lower sample amounts may impact results. Please discuss with your Metabolon Client Success Project Coordinator.
- Additional volume or mass may be required for a longitudinal bridging matrix. Please discuss with your Metabolon Client Success Project Coordinator.

Unique Sample Types Guidance

 Sample types not listed in the above table may require specific instructions. Please discuss with your Metabolon Client Success Project Coordinator

Assay-Specific Considerations

Targeted Assays and Metal Ions Panel

Sample amounts for these specific assays may vary. Please discuss with your specific Metabolon Project Coordinator.